

Generalized Fibonacci numbers and dimer statistics

W. T. Lu and F. Y. Wu

Department of Physics, Northeastern University, Boston, Massachusetts 02115

Abstract

We establish new product identities involving the q -analogue of the Fibonacci numbers. We show that the identities lead to alternate expressions of generating functions for close-packed dimers on non-orientable surfaces.

The recursion relation

$$\mathcal{F}_{n+2} = \mathcal{F}_n + \mathcal{F}_{n+1} \quad (1)$$

with the initial values $\mathcal{F}_0 = \mathcal{F}_1 = 1$ produces the well-known Fibonacci (1170-1250) numbers 1, 1, 2, 3, 5, 8, 13, 21, 34, 55, $\dots$. The Fibonacci sequence can be generalized in a number of ways [1, 2]. One generalization is the q -analogue of the sequence defined by the recursion relation

$$\mathcal{F}_{n+2}(q) = \mathcal{F}_n(q) + q\mathcal{F}_{n+1}(q), \quad (2)$$

and the initial condition $\mathcal{F}_0(q) = 1$ and $\mathcal{F}_1(q) = q$. Here q is any real or complex number. This yields the sequence 1, q , $q^2 + 1$, $q^3 + 2q$, $q^4 + 3q^2 + 1$, $\dots$.

It can be readily established that the generating function of $\mathcal{F}_n(q)$ is

$$\frac{1}{1 - qs - s^2} = \sum_{n=0}^{\infty} \mathcal{F}_n(q) s^n. \quad (3)$$

Writing $q = x - x^{-1}$ and partial fractioning the left-hand side of (3), one obtains the explicit expression

$$\mathcal{F}_n(q) = \frac{x^{n+1} + (-1)^n x^{-(n+1)}}{x + x^{-1}}, \quad n = 0, 1, 2, \dots \quad (4)$$

We have following identities which we state as a theorem.

Theorem: (A) For $M = \text{integers}$ and $\theta_m = m\pi/(2M + 1)$, we have

$$\mathcal{F}_M(q) \pm i \mathcal{F}_{M-1}(q) = \prod_{m=1}^M \left(q \mp 2i (-1)^m \cos \theta_m \right), \quad M = 1, 2, \dots \quad (5)$$

(B) For $M = \text{integers}$ we have

$$\mathcal{F}_M(q) = \prod_{m=1}^M \left(q + 2i \cos \frac{m\pi}{M+1} \right), \quad M = 1, 2, \dots \quad (6)$$

(C) For $N = \text{integers}$ and $\phi_n = (2n - 1)\pi/2N$, we have

$$\mathcal{F}_N(q) + \mathcal{F}_{N-2}(q) = \prod_{n=1}^N \left(q \pm 2i \cos \phi_n \right) \quad N = 2, 4, \dots \quad (7)$$

$$= \pm 2i + \prod_{n=1}^N \left(q \pm 2i (-1)^n \sin \phi_n \right), \quad N = 1, 3, \dots \quad (8)$$

(D) For $M = \text{integers}$, $N = \text{odd}$, $q_n = 2\tau^{-1} \sin \phi_n$, and $p_m = 2\tau \cos \theta_m$, we have

$$\tau^{MN} \prod_{n=1}^N \left(\mathcal{F}_M(q_n) - i (-1)^{n+M} \mathcal{F}_{M-1}(q_n) \right) = i^{-M^2} \prod_{m=1}^M \left(\mathcal{F}_N(p_m) + \mathcal{F}_{N-2}(p_m) + 2i(-1)^{m+M} \right), \quad (9)$$

These identities can be established by determining the zeroes of the expressions. To establish (A), for example, using (4) it is straightforward to deduce the identity

$$\mathcal{F}_M(q) \pm i \mathcal{F}_{M-1}(q) = x^{-M} \left(\frac{x^{2M+1} - (\pm i)^{2M+1}}{x \mp i} \right). \quad (10)$$

The quantity inside the parentheses in (10) is a polynomial of degree $2M$ in x with the highest term x^{2M} . It has $2M$ zeroes at

$$x_m = \pm i e^{i\theta_{2m}}, \quad m = 1, 2, \dots, 2M.$$

Using the fact that $x_{2M} = -x_1^*$, $x_{2M-1} = -x_2^*$, ..., $x_{M+1} = -x_M^*$, we obtain

$$\begin{aligned} \mathcal{F}_M(q) \pm i \mathcal{F}_{M-1}(q) &= x^{-M} \prod_{m=1}^{2M} (x - x_m) \\ &= x^{-M} \prod_{m=1}^M (x - x_m)(x + x_m^*) \\ &= \prod_{m=1}^M \left(q \mp 2i \cos \theta_{2m} \right). \end{aligned} \quad (11)$$

This leads to (A) after we make use of the identities $\cos \theta_{2M} = -\cos \theta_1$, $\cos \theta_{2M-2} = (-1)^3 \cos \theta_3, \dots$. The identity (B) is proved in a similar fashion.

To establish (C) we note that for all N we have

$$\mathcal{F}_N(q) + \mathcal{F}_{N-2}(q) = x^N + (-x)^{-N}.$$

For $N = \text{even}$, the polynomial $x^{-N}(x^{2N} + 1)$ has $2N$ zeroes at

$$x_n = \mp i e^{i\phi_n}, \quad x_n = \mp i e^{-i\phi_n}, \quad n = 1, 2, \dots, N. \quad (12)$$

This leads to (7) by factorizing the polynomial as in the proof of (A).

For $N = \text{odd}$, we have instead

$$\begin{aligned} \mathcal{F}_N(q) + \mathcal{F}_{N-2}(q) \mp 2i &= x^N - x^{-N} \mp 2i \\ &= x^{-N} \left(x^N \mp i \right)^2 \\ &= x^{-N} \prod_{n=1}^N (x - x_n)^2 \end{aligned} \quad (13)$$

where $x_n = \mp (-1)^n e^{i\phi_n}$. The identity (8) now follows the replacement of $x_1 = -x_N^*$, $x_2 = -x_{N-1}^*, \dots$ in one of the $x - x_n$ factors.

To establish (D), write $q = q_n = 2\tau^{-1} \sin \phi_n$ in (A) and $q = p_m = 2\tau \cos \theta_m$ in (C) to obtain, respectively,

$$\begin{aligned}\mathcal{F}_M(q_n) - i(-1)^{n+M} \mathcal{F}_{M-1}(q_n) &= \prod_{m=1}^M \left(2\tau^{-1} \sin \phi_n + 2i(-1)^{m+n+M} \cos \theta_m \right) \\ \mathcal{F}_N(p_m) + \mathcal{F}_{N-2}(p_m) + 2i(-1)^{m+M} &= \prod_{n=1}^N \left(2\tau \cos \theta_m - 2i(-1)^{n+m+M} \sin \phi_n \right).\end{aligned}$$

The identity (D) now follows by combining these two expressions and the fact that, for $N = \text{odd}$ integers,

$$\prod_{m=1}^M \prod_{n=1}^N \left(i(-1)^{m+M+n+1} \right) = i^{M^2}.$$

The identities (A) - (D) can be used to deduce alternate expressions for dimer generating functions on a Möbius strip. Let $Z_{\mathcal{M}, \mathcal{N}}^{\text{Mob}}(z_h, z_v)$ be the generating function on an $\mathcal{M} \times \mathcal{N}$ Möbius strip of width $\mathcal{N}$ with horizontal dimer weights z_h and vertical dimer weights z_v . It can be shown that, by using (A) - (D) and the equivalences of the sets $\{\sin \phi_{2n}\} = \{\cos \phi_n\}$ for $N = \text{even}$ and $\{\sin \phi_{2n}\} = \{(-1)^n \sin \phi_n\}$ with $n = 1, 2, \dots, N$, one establishes the following equivalent expressions:

For $N = \text{even}$ and $\tau = z_v/z_h$,

$$Z_{2M, N}^{\text{Mob}}(z_h, z_v) = \prod_{m=1}^M \prod_{n=1}^{N/2} \left(4z_h^2 \sin^2 \phi_{2n} + 4z_v^2 \cos^2 \theta_m \right) \quad (14)$$

$$= z_h^{MN} \prod_{m=1}^M \left(\mathcal{F}_N(p_m) + \mathcal{F}_{N-2}(p_m) \right) \quad (15)$$

$$= z_v^{MN} \prod_{n=1}^{N/2} \mathcal{F}_{2M}(q_n), \quad N = \text{even} \quad (16)$$

$$Z_{2M-1, N}^{\text{Mob}}(z_h, z_v) = 2z_h^{N/2} \prod_{m=1}^{M-1} \prod_{n=1}^{N/2} \left(4z_h^2 \sin^2 \phi_{2n} + 4z_v^2 \cos^2(m\pi/2M) \right) \quad (17)$$

$$= 2z_h^{(2M-1)N/2} \prod_{m=1}^{M-1} \left[\mathcal{F}_N(2\tau \cos(m\pi/2M)) + \mathcal{F}_{N-2}(2\tau \cos(m\pi/2M)) \right] \quad (18)$$

$$= 2z_h^{N/2} z_v^{(M-1)N} \prod_{n=1}^{N/2} \left(q_n^{-1} \mathcal{F}_{2M-1}(q_n) \right), \quad N = \text{even}. \quad (19)$$

For $N = \text{odd}$ and $\tau = z_v/z_h$,

$$Z_{2M, N}^{\text{Mob}}(z_h, z_v) = \text{Re} \left[(1-i) \prod_{m=1}^M \prod_{n=1}^N \left(2i(-1)^{M+m+1} z_h \sin \phi_{2n} + 2z_v \cos \theta_m \right) \right] \quad (20)$$

$$= z_h^{MN} \text{Re} \left[(1-i) \prod_{m=1}^M \left(\mathcal{F}_N(p_m) + \mathcal{F}_{N-2}(p_m) + 2i(-1)^{m+M} \right) \right] \quad (21)$$

$$= z_v^{MN} \text{Re} \left[(1-i) i^{M^2} \prod_{n=1}^N \left(\mathcal{F}_M(q_n) - i(-1)^{n+M} \mathcal{F}_{M-1}(q_n) \right) \right], \quad N = \text{odd}. \quad (22)$$

Expressions (14), (17) and (20) are those given by Lu and Wu [3, 4], expressions (15), (18) and (21) are given by Tesler [5], and expressions (16), (19) and (22) are new.

Work has been supported in part by NSF grant DMR-9980440.

References

- [1] G. E. Andrews, *Number Theory* (W. B. Saunders, Philadelphia, 1971 and Dover, New York, 1994) p. 23.
- [2] S. W. Golomb and A. Lempel, Second order polynomial recursions, SIAM J. Appl. Math. **33**, 587-592 (1977)
- [3] W. T. Lu and F. Y. Wu, Dimer statistics on non-orientable surfaces, Phys. Lett. A **259**, 108-114 (1999).
- [4] W. T. Lu and F. Y. Wu, Close-packed dimers on non-orientable surfaces, Phys. Lett. A **293**, 235-246 (2002); Erratum, *ibid.* **298**, 293 (2002).
- [5] G. Tesler, Perfect matchings in graphs on non-orientable surfaces, J. Comb. Theo. B **78**, 198-231 (2000).